

6. 呼吸器

(1) 胸部X線検査

① 1日人間ドック

<胸部X線検査判定結果>

<男性>

| | ~29 | 30~39 | 40~49 | 50~59 | 60~69 | 70~79 | 80~ | 合計 (人) | 比率 (%) |
|----------|-----|-------|-------|-------|-------|-------|------|-----------|-----------|
| A | 33 | 1,244 | 3,563 | 3,537 | 2,140 | 416 | 41 | 10,974 | 76.3 |
| B1 | 1 | 84 | 440 | 729 | 763 | 305 | 66 | 2,388 | 16.6 |
| B2 | 0 | 28 | 113 | 238 | 266 | 138 | 41 | 824 | 5.7 |
| C | 0 | 0 | 2 | 2 | 2 | 1 | 0 | 7 | 0.0 |
| D | 0 | 4 | 30 | 52 | 68 | 36 | 6 | 196 | 1.4 |
| E | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0.0 |
| 受診者数(人) | 34 | 1,360 | 4,148 | 4,558 | 3,239 | 896 | 154 | 14,389 | 100.0 |
| 有所見者数(人) | 1 | 116 | 585 | 1,021 | 1,099 | 480 | 113 | 3,415 | 23.7 |
| 有所見者率(%) | 2.9 | 8.5 | 14.1 | 22.4 | 33.9 | 53.6 | 73.4 | 23.7 | |

<女性>

| | ~29 | 30~39 | 40~49 | 50~59 | 60~69 | 70~79 | 80~ | 合計 (人) | 比率 (%) |
|----------|-----|-------|-------|-------|-------|-------|------|-----------|-----------|
| A | 34 | 1,132 | 3,123 | 2,959 | 1,453 | 324 | 25 | 9,050 | 78.9 |
| B1 | 3 | 60 | 317 | 570 | 648 | 251 | 45 | 1,894 | 16.5 |
| B2 | 0 | 13 | 49 | 110 | 135 | 60 | 18 | 385 | 3.4 |
| C | 0 | 0 | 2 | 1 | 4 | 2 | 0 | 9 | 0.1 |
| D | 0 | 6 | 26 | 34 | 46 | 18 | 3 | 133 | 1.2 |
| E | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0.0 |
| 受診者数(人) | 37 | 1,211 | 3,517 | 3,674 | 2,286 | 655 | 91 | 11,471 | 100.0 |
| 有所見者数(人) | 3 | 79 | 394 | 715 | 833 | 331 | 66 | 2,421 | 21.1 |
| 有所見者率(%) | 8.1 | 6.5 | 11.2 | 19.5 | 36.4 | 50.5 | 72.5 | 21.1 | |

<胸部X線検査疾患別一覧>

<男性>

| | ～29 | 30～39 | 40～49 | 50～59 | 60～69 | 70～79 | 80～ | 総計 |
|---------|-----|-------|-------|-------|-------|-------|-----|--------|
| 著変なし | 33 | 1,244 | 3,563 | 3,537 | 2,140 | 416 | 41 | 10,974 |
| 硬化性陰影 | 0 | 34 | 172 | 270 | 302 | 149 | 37 | 964 |
| 線状影 | 0 | 15 | 55 | 97 | 105 | 42 | 8 | 322 |
| 索状影 | 0 | 2 | 26 | 48 | 66 | 39 | 13 | 194 |
| 気腫性変化 | 0 | 15 | 70 | 155 | 143 | 34 | 6 | 423 |
| 限局性陰影 | 0 | 4 | 20 | 32 | 38 | 22 | 1 | 117 |
| 網状影 | 0 | 1 | 1 | 27 | 60 | 42 | 23 | 154 |
| 結節影 | 0 | 2 | 22 | 28 | 33 | 16 | 4 | 105 |
| 粒状影 | 0 | 3 | 4 | 7 | 12 | 4 | 1 | 31 |
| 浸潤影 | 0 | 0 | 1 | 6 | 3 | 3 | 0 | 13 |
| 縦隔・肺門増大 | 0 | 0 | 1 | 9 | 2 | 2 | 2 | 16 |
| 腫瘤影 | 0 | 1 | 1 | 3 | 3 | 1 | 0 | 9 |
| 胸水 | 0 | 1 | 1 | 0 | 1 | 2 | 0 | 5 |
| 気胸 | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 1 |
| 受診者数(人) | 34 | 1,360 | 4,148 | 4,558 | 3,239 | 896 | 154 | 14,389 |

<女性>

| | ～29 | 30～39 | 40～49 | 50～59 | 60～69 | 70～79 | 80～ | 総計 |
|---------|-----|-------|-------|-------|-------|-------|-----|--------|
| 著変なし | 34 | 1,132 | 3,123 | 2,959 | 1,453 | 324 | 25 | 9,050 |
| 硬化性陰影 | 1 | 27 | 154 | 286 | 298 | 120 | 24 | 910 |
| 線状影 | 0 | 11 | 35 | 50 | 60 | 22 | 5 | 183 |
| 索状影 | 1 | 5 | 18 | 42 | 55 | 21 | 1 | 143 |
| 気腫性変化 | 0 | 1 | 9 | 14 | 23 | 10 | 2 | 59 |
| 限局性陰影 | 0 | 3 | 15 | 20 | 28 | 7 | 3 | 76 |
| 網状影 | 0 | 1 | 2 | 5 | 9 | 9 | 6 | 32 |
| 結節影 | 0 | 3 | 13 | 18 | 26 | 10 | 0 | 70 |
| 粒状影 | 0 | 0 | 5 | 5 | 10 | 3 | 0 | 23 |
| 浸潤影 | 0 | 0 | 3 | 4 | 6 | 1 | 0 | 14 |
| 縦隔・肺門増大 | 0 | 1 | 3 | 1 | 1 | 0 | 0 | 6 |
| 腫瘤影 | 0 | 0 | 0 | 2 | 0 | 1 | 0 | 3 |
| 胸水 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 気胸 | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 1 |
| 受診者数(人) | 37 | 1,211 | 3,517 | 3,674 | 2,286 | 655 | 91 | 11,471 |

② 2日人間ドック

<胸部X線検査判定結果>

<男性>

| | ~29 | 30~39 | 40~49 | 50~59 | 60~69 | 70~79 | 80~ | 合計 (人) | 比率 (%) |
|----------|------|-------|-------|-------|-------|-------|------|-----------|-----------|
| A | 7 | 212 | 582 | 834 | 791 | 313 | 68 | 2,807 | 74.8 |
| B1 | 1 | 13 | 45 | 140 | 210 | 170 | 68 | 647 | 17.2 |
| B2 | 0 | 4 | 29 | 52 | 87 | 48 | 33 | 253 | 6.7 |
| C | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 1 | 0.0 |
| D | 0 | 0 | 4 | 9 | 15 | 7 | 10 | 45 | 1.2 |
| E | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0.0 |
| 受診者数(人) | 8 | 229 | 660 | 1,035 | 1,103 | 539 | 179 | 3,753 | 100.0 |
| 有所見者数(人) | 1 | 17 | 78 | 201 | 312 | 226 | 111 | 946 | 25.2 |
| 有所見者率(%) | 12.5 | 7.4 | 11.8 | 19.4 | 28.3 | 41.9 | 62.0 | 25.2 | |

<女性>

| | ~29 | 30~39 | 40~49 | 50~59 | 60~69 | 70~79 | 80~ | 合計 (人) | 比率 (%) |
|----------|-----|-------|-------|-------|-------|-------|------|-----------|-----------|
| A | 3 | 58 | 169 | 319 | 377 | 153 | 24 | 1,103 | 72.9 |
| B1 | 0 | 1 | 11 | 43 | 117 | 110 | 38 | 320 | 21.2 |
| B2 | 0 | 2 | 2 | 10 | 22 | 27 | 14 | 77 | 5.1 |
| C | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 1 | 0.1 |
| D | 0 | 1 | 0 | 2 | 4 | 5 | 0 | 12 | 0.8 |
| E | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0.0 |
| 受診者数(人) | 3 | 62 | 183 | 374 | 520 | 295 | 76 | 1,513 | 100.0 |
| 有所見者数(人) | 0 | 4 | 14 | 55 | 143 | 142 | 52 | 410 | 27.1 |
| 有所見者率(%) | 0.0 | 6.5 | 7.7 | 14.7 | 27.5 | 48.1 | 68.4 | 27.1 | |

<胸部X線検査疾患別一覧>

<男性>

| | ～29 | 30～39 | 40～49 | 50～59 | 60～69 | 70～79 | 80～ | 総計 |
|---------|-----|-------|-------|-------|-------|-------|-----|-------|
| 著変なし | 7 | 212 | 582 | 834 | 791 | 313 | 68 | 2,807 |
| 硬化性陰影 | 0 | 3 | 16 | 49 | 89 | 74 | 44 | 275 |
| 線状影 | 0 | 2 | 13 | 18 | 29 | 24 | 7 | 93 |
| 索状影 | 0 | 2 | 2 | 14 | 26 | 16 | 11 | 71 |
| 気腫性変化 | 1 | 3 | 25 | 61 | 67 | 32 | 20 | 209 |
| 限局性陰影 | 0 | 1 | 4 | 9 | 14 | 4 | 2 | 34 |
| 網状影 | 0 | 0 | 1 | 3 | 17 | 22 | 15 | 58 |
| 結節影 | 0 | 1 | 1 | 5 | 7 | 2 | 2 | 18 |
| 粒状影 | 0 | 0 | 2 | 0 | 2 | 4 | 1 | 9 |
| 浸潤影 | 0 | 0 | 0 | 0 | 3 | 2 | 1 | 6 |
| 縦隔・肺門増大 | 0 | 0 | 0 | 0 | 1 | 1 | | 2 |
| 腫瘤影 | 0 | 0 | 0 | 2 | 2 | 0 | 2 | 6 |
| 胸水 | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 1 |
| 気胸 | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 1 |
| 受診者数(人) | 8 | 229 | 660 | 1,035 | 1,103 | 539 | 179 | 3,753 |

<女性>

| | ～29 | 30～39 | 40～49 | 50～59 | 60～69 | 70～79 | 80～ | 総計 |
|---------|-----|-------|-------|-------|-------|-------|-----|-------|
| 著変なし | 3 | 58 | 169 | 319 | 377 | 153 | 24 | 1,103 |
| 硬化性陰影 | 0 | 1 | 4 | 22 | 49 | 39 | 23 | 138 |
| 線状影 | 0 | 0 | 3 | 5 | 8 | 13 | 3 | 32 |
| 索状影 | 0 | 0 | 0 | 1 | 20 | 18 | 4 | 43 |
| 気腫性変化 | 0 | 0 | 1 | 3 | 10 | 7 | 3 | 24 |
| 限局性陰影 | 0 | 1 | 1 | 3 | 4 | 4 | 0 | 13 |
| 網状影 | 0 | 1 | 0 | 1 | 4 | 2 | 4 | 12 |
| 結節影 | 0 | 0 | 0 | 0 | 2 | 2 | 0 | 4 |
| 粒状影 | 0 | 0 | 0 | 0 | 1 | 2 | 0 | 3 |
| 浸潤影 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 縦隔・肺門増大 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 1 |
| 腫瘤影 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 1 |
| 胸水 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 気胸 | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 1 |
| 受診者数(人) | 3 | 62 | 183 | 374 | 520 | 295 | 76 | 1,513 |

③ 施設内健診（協会けんぽ含む）

<胸部 X 線検査判定結果>

<男性>

| | ～29 | 30～39 | 40～49 | 50～59 | 60～69 | 70～79 | 80～ | 合計 (人) | 比率 (%) |
|-----------|-------|-------|-------|-------|-------|-------|------|-----------|-----------|
| A | 1,892 | 3,273 | 3,221 | 2,367 | 1,530 | 102 | 4 | 12,389 | 86.1 |
| B1 | 45 | 148 | 280 | 333 | 361 | 38 | 1 | 1,206 | 8.4 |
| B2 | 21 | 77 | 106 | 166 | 221 | 50 | 1 | 642 | 4.5 |
| C | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 1 | 0.0 |
| D | 8 | 20 | 27 | 40 | 64 | 0 | 0 | 159 | 1.1 |
| E | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0.0 |
| 受診者数 (人) | 1,966 | 3,518 | 3,634 | 2,907 | 2,176 | 190 | 6 | 14,397 | 100.0 |
| 有所見者数 (人) | 74 | 245 | 413 | 540 | 646 | 88 | 2 | 2,008 | 13.9 |
| 有所見者率 (%) | 3.8 | 7.0 | 11.4 | 18.6 | 29.7 | 46.3 | 33.3 | 13.9 | |

<女性>

| | ～29 | 30～39 | 40～49 | 50～59 | 60～69 | 70～79 | 80～ | 合計 (人) | 比率 (%) |
|-----------|-------|-------|-------|-------|-------|-------|------|-----------|-----------|
| A | 2,332 | 3,044 | 3,525 | 2,585 | 891 | 50 | 7 | 12,434 | 89.9 |
| B1 | 52 | 82 | 218 | 326 | 213 | 27 | 2 | 920 | 6.7 |
| B2 | 7 | 24 | 56 | 120 | 97 | 8 | 3 | 315 | 2.3 |
| C | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0.0 |
| D | 6 | 22 | 30 | 59 | 39 | 9 | 0 | 165 | 1.2 |
| E | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0.0 |
| 受診者数 (人) | 2,397 | 3,172 | 3,829 | 3,090 | 1,240 | 94 | 12 | 13,834 | 100.0 |
| 有所見者数 (人) | 65 | 128 | 304 | 505 | 349 | 44 | 5 | 1,400 | 10.1 |
| 有所見者率 (%) | 2.7 | 4.0 | 7.9 | 16.3 | 28.1 | 46.8 | 41.7 | 10.1 | |

<胸部X線検査疾患別一覧>

<男性>

| | ～29 | 30～39 | 40～49 | 50～59 | 60～69 | 70～79 | 80～ | 総計 |
|---------|-------|-------|-------|-------|-------|-------|-----|--------|
| 著変なし | 1,892 | 3,273 | 3,221 | 2,367 | 1,530 | 102 | 4 | 12,389 |
| 硬化性陰影 | 18 | 52 | 129 | 161 | 181 | 29 | 1 | 571 |
| 線状影 | 6 | 23 | 38 | 60 | 75 | 5 | 0 | 207 |
| 索状影 | 3 | 25 | 33 | 49 | 59 | 8 | 0 | 177 |
| 気腫性変化 | 4 | 45 | 86 | 118 | 110 | 13 | 0 | 376 |
| 限局性陰影 | 3 | 11 | 14 | 17 | 29 | 0 | 0 | 74 |
| 網状影 | 0 | 4 | 4 | 16 | 52 | 12 | 1 | 89 |
| 結節影 | 2 | 5 | 8 | 15 | 22 | 1 | 0 | 53 |
| 粒状影 | 0 | 5 | 5 | 7 | 10 | 2 | 0 | 29 |
| 浸潤影 | 0 | 2 | 1 | 4 | 6 | 0 | 0 | 13 |
| 縦隔・肺門増大 | 1 | 3 | 0 | 5 | 4 | 0 | 0 | 13 |
| 腫瘤影 | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 1 |
| 胸水 | 0 | 1 | 1 | 0 | 3 | 0 | 0 | 5 |
| 気胸 | 1 | 0 | 0 | 1 | 0 | 0 | 0 | 2 |
| 受診者数(人) | 1,966 | 3,518 | 3,634 | 2,907 | 2,176 | 190 | 6 | 14,397 |

<女性>

| | ～29 | 30～39 | 40～49 | 50～59 | 60～69 | 70～79 | 80～ | 総計 |
|---------|-------|-------|-------|-------|-------|-------|-----|--------|
| 著変なし | 2,332 | 3,044 | 3,525 | 2,585 | 891 | 50 | 7 | 12,434 |
| 硬化性陰影 | 12 | 53 | 123 | 221 | 123 | 13 | 1 | 546 |
| 線状影 | 7 | 14 | 28 | 57 | 29 | 6 | 1 | 142 |
| 索状影 | 2 | 5 | 8 | 36 | 31 | 6 | 2 | 90 |
| 気腫性変化 | 0 | 2 | 10 | 13 | 15 | 0 | 0 | 40 |
| 限局性陰影 | 2 | 6 | 10 | 21 | 17 | 5 | 0 | 61 |
| 網状影 | 0 | 0 | 1 | 10 | 3 | 4 | 0 | 18 |
| 結節影 | 2 | 11 | 13 | 29 | 14 | 4 | 0 | 73 |
| 粒状影 | 0 | 1 | 3 | 13 | 11 | 2 | 0 | 30 |
| 浸潤影 | 0 | 1 | 7 | 7 | 5 | 0 | 0 | 20 |
| 縦隔・肺門増大 | 0 | 0 | 5 | 3 | 2 | 0 | 0 | 10 |
| 腫瘤影 | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 1 |
| 胸水 | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 1 |
| 気胸 | 1 | 0 | 1 | 0 | 0 | 0 | 0 | 2 |
| 受診者数(人) | 2,397 | 3,172 | 3,829 | 3,090 | 1,240 | 94 | 12 | 13,834 |

④ 施設外健診（協会けんぽ含む）

<胸部 X 線検査判定結果>

<男性>

| | ～29 | 30～39 | 40～49 | 50～59 | 60～69 | 70～79 | 80～ | 合計 (人) | 比率 (%) |
|-----------|-------|-------|-------|-------|-------|-------|------|-----------|-----------|
| A | 4,202 | 4,350 | 3,559 | 3,375 | 2,695 | 1,119 | 369 | 19,669 | 86.1 |
| B1 | 71 | 182 | 300 | 441 | 519 | 390 | 210 | 2,113 | 9.2 |
| B2 | 10 | 42 | 83 | 141 | 208 | 187 | 143 | 814 | 3.6 |
| C | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 1 | 0.0 |
| D | 6 | 16 | 31 | 43 | 76 | 55 | 28 | 255 | 1.1 |
| E | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0.0 |
| 受診者数 (人) | 4,289 | 4,590 | 3,974 | 4,000 | 3,498 | 1,751 | 750 | 22,852 | 100.0 |
| 有所見者数 (人) | 87 | 240 | 415 | 625 | 803 | 632 | 381 | 3,183 | 13.9 |
| 有所見者率 (%) | 2.0 | 5.2 | 10.4 | 15.6 | 23.0 | 36.1 | 50.8 | 13.9 | |

<女性>

| | ～29 | 30～39 | 40～49 | 50～59 | 60～69 | 70～79 | 80～ | 合計 (人) | 比率 (%) |
|-----------|-------|-------|-------|-------|-------|-------|------|-----------|-----------|
| A | 2,161 | 1,780 | 1,904 | 1,943 | 2,243 | 1,549 | 456 | 12,036 | 84.2 |
| B1 | 34 | 42 | 96 | 233 | 395 | 507 | 267 | 1,574 | 11.0 |
| B2 | 7 | 7 | 23 | 67 | 120 | 157 | 128 | 509 | 3.6 |
| C | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0.0 |
| D | 2 | 5 | 10 | 33 | 46 | 52 | 24 | 172 | 1.2 |
| E | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0.0 |
| 受診者数 (人) | 2,204 | 1,834 | 2,033 | 2,276 | 2,804 | 2,265 | 875 | 14,291 | 100.0 |
| 有所見者数 (人) | 43 | 54 | 129 | 333 | 561 | 716 | 419 | 2,255 | 15.8 |
| 有所見者率 (%) | 2.0 | 2.9 | 6.3 | 14.6 | 20.0 | 31.6 | 47.9 | 15.8 | |

<胸部X線検査疾患別一覧>

<男性>

| | ～29 | 30～39 | 40～49 | 50～59 | 60～69 | 70～79 | 80～ | 総計 |
|---------|-------|-------|-------|-------|-------|-------|-----|--------|
| 著変なし | 4,202 | 4,350 | 3,559 | 3,375 | 2,695 | 1,119 | 369 | 19,669 |
| 硬化性陰影 | 19 | 89 | 130 | 204 | 262 | 194 | 122 | 1,020 |
| 線状影 | 13 | 36 | 52 | 78 | 83 | 80 | 54 | 396 |
| 索状影 | 5 | 6 | 18 | 29 | 32 | 35 | 15 | 140 |
| 気腫性変化 | 8 | 28 | 92 | 126 | 138 | 73 | 35 | 500 |
| 限局性陰影 | 2 | 14 | 19 | 31 | 46 | 33 | 17 | 162 |
| 網状影 | 1 | 2 | 3 | 23 | 46 | 73 | 68 | 216 |
| 結節影 | 2 | 12 | 16 | 24 | 30 | 22 | 11 | 117 |
| 粒状影 | 0 | 1 | 3 | 5 | 7 | 5 | 3 | 24 |
| 浸潤影 | 1 | 2 | 2 | 1 | 6 | 10 | 3 | 25 |
| 縦隔・肺門増大 | 1 | 1 | 0 | 5 | 4 | 4 | 6 | 21 |
| 腫瘤影 | 0 | 1 | 1 | 2 | 2 | 2 | 0 | 8 |
| 胸水 | 0 | 1 | 0 | 1 | 3 | 4 | 2 | 11 |
| 気胸 | 2 | 2 | 0 | 0 | 0 | 0 | 0 | 4 |
| 受診者数(人) | 4,289 | 4,590 | 3,974 | 4,000 | 3,498 | 1,751 | 750 | 22,852 |

<女性>

| | ～29 | 30～39 | 40～49 | 50～59 | 60～69 | 70～79 | 80～ | 総計 |
|---------|-------|-------|-------|-------|-------|-------|-----|--------|
| 著変なし | 2,161 | 1,780 | 1,904 | 1,943 | 2,243 | 1,549 | 456 | 12,036 |
| 硬化性陰影 | 8 | 11 | 54 | 137 | 204 | 202 | 112 | 728 |
| 線状影 | 2 | 6 | 13 | 34 | 98 | 104 | 56 | 313 |
| 索状影 | 1 | 4 | 3 | 11 | 26 | 45 | 20 | 110 |
| 気腫性変化 | 0 | 3 | 5 | 8 | 9 | 11 | 7 | 43 |
| 限局性陰影 | 0 | 3 | 6 | 23 | 37 | 39 | 22 | 130 |
| 網状影 | 1 | 1 | 1 | 5 | 16 | 26 | 38 | 88 |
| 結節影 | 2 | 2 | 3 | 19 | 20 | 23 | 9 | 78 |
| 粒状影 | 0 | 1 | 1 | 5 | 8 | 5 | 2 | 22 |
| 浸潤影 | 1 | 1 | 5 | 9 | 6 | 7 | 5 | 34 |
| 縦隔・肺門増大 | 1 | 0 | 0 | 0 | 3 | 3 | 4 | 11 |
| 腫瘤影 | 0 | 0 | 0 | 0 | 1 | 0 | 1 | 2 |
| 胸水 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 1 |
| 気胸 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 受診者数(人) | 2,204 | 1,834 | 2,033 | 2,276 | 2,804 | 2,265 | 875 | 14,291 |

⑤ 全国健康保険協会管掌健康保険生活習慣病予防健診[再掲]

<胸部 X 線検査判定結果>

<男性>

| | ~29 | 30~39 | 40~49 | 50~59 | 60~69 | 70~79 | 80~ | 合計 (人) | 比率 (%) |
|-----------|-----|-------|-------|-------|-------|-------|-----|-----------|-----------|
| A | 0 | 1,855 | 3,025 | 2,243 | 1,423 | 80 | 0 | 8,626 | 83.1 |
| B1 | 0 | 94 | 275 | 339 | 330 | 29 | 0 | 1,067 | 10.3 |
| B2 | 0 | 48 | 97 | 167 | 194 | 39 | 0 | 545 | 5.3 |
| C | 0 | 0 | 1 | 1 | 0 | 0 | 0 | 2 | 0.0 |
| D | 0 | 9 | 30 | 39 | 59 | 0 | 0 | 137 | 1.3 |
| E | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0.0 |
| 受診者数 (人) | 0 | 2,006 | 3,428 | 2,789 | 2,006 | 148 | 0 | 10,377 | 100.0 |
| 有所見者数 (人) | 0 | 151 | 403 | 546 | 583 | 68 | 0 | 1,751 | 16.9 |
| 有所見者率 (%) | 0.0 | 7.5 | 11.8 | 19.6 | 29.1 | 45.9 | 0.0 | 16.9 | |

<女性>

| | ~29 | 30~39 | 40~49 | 50~59 | 60~69 | 70~79 | 80~ | 合計 (人) | 比率 (%) |
|-----------|-----|-------|-------|-------|-------|-------|-----|-----------|-----------|
| A | 0 | 1,129 | 2,196 | 1,582 | 642 | 32 | 0 | 5,581 | 87.1 |
| B1 | 0 | 26 | 132 | 222 | 148 | 14 | 0 | 542 | 8.5 |
| B2 | 0 | 11 | 45 | 74 | 69 | 3 | 0 | 202 | 3.2 |
| C | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0.0 |
| D | 0 | 8 | 14 | 34 | 25 | 1 | 0 | 82 | 1.3 |
| E | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0.0 |
| 受診者数 (人) | 0 | 1,174 | 2,387 | 1,912 | 884 | 50 | 0 | 6,407 | 100.0 |
| 有所見者数 (人) | 0 | 45 | 191 | 330 | 242 | 18 | 0 | 826 | 12.9 |
| 有所見者率 (%) | 0.0 | 3.8 | 8.0 | 17.3 | 27.4 | 36.0 | 0.0 | 12.9 | |

<胸部X線検査疾患別一覧>

<男性>

| | ～29 | 30～39 | 40～49 | 50～59 | 60～69 | 70～79 | 80～ | 総計 |
|---------|-----|-------|-------|-------|-------|-------|-----|--------|
| 著変なし | 0 | 1,855 | 3,025 | 2,243 | 1,423 | 80 | 0 | 8,626 |
| 硬化性陰影 | 0 | 41 | 110 | 158 | 173 | 21 | 0 | 503 |
| 線状影 | 0 | 13 | 39 | 63 | 67 | 4 | 0 | 186 |
| 索状影 | 0 | 12 | 27 | 51 | 48 | 6 | 0 | 144 |
| 気腫性変化 | 0 | 28 | 104 | 122 | 92 | 10 | 0 | 356 |
| 限局性陰影 | 0 | 6 | 18 | 20 | 27 | 0 | 0 | 71 |
| 網状影 | 0 | 2 | 4 | 19 | 50 | 10 | 0 | 85 |
| 結節影 | 0 | 4 | 9 | 21 | 17 | 1 | 0 | 52 |
| 粒状影 | 0 | 2 | 4 | 6 | 8 | 2 | 0 | 22 |
| 浸潤影 | 0 | 1 | 2 | 4 | 6 | 0 | 0 | 13 |
| 縦隔・肺門増大 | 0 | 2 | 0 | 4 | 4 | 0 | 0 | 10 |
| 腫瘤影 | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 1 |
| 胸水 | 0 | 0 | 1 | 0 | 3 | 0 | 0 | 4 |
| 気胸 | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 1 |
| 受診者数(人) | 0 | 2,006 | 3,428 | 2,789 | 2,006 | 148 | 0 | 10,377 |

<女性>

| | ～29 | 30～39 | 40～49 | 50～59 | 60～69 | 70～79 | 80～ | 総計 |
|---------|-----|-------|-------|-------|-------|-------|-----|-------|
| 著変なし | 0 | 1,129 | 2,196 | 1,582 | 642 | 32 | 0 | 5,581 |
| 硬化性陰影 | 0 | 16 | 85 | 141 | 83 | 7 | 0 | 332 |
| 線状影 | 0 | 8 | 16 | 39 | 17 | 3 | 0 | 83 |
| 索状影 | 0 | 5 | 8 | 25 | 25 | 3 | 0 | 66 |
| 気腫性変化 | 0 | 1 | 7 | 8 | 10 | 0 | 0 | 26 |
| 限局性陰影 | 0 | 2 | 4 | 12 | 12 | 1 | 0 | 31 |
| 網状影 | 0 | 0 | 1 | 4 | 3 | 1 | 0 | 9 |
| 結節影 | 0 | 4 | 9 | 17 | 13 | 1 | 0 | 44 |
| 粒状影 | 0 | 0 | 2 | 7 | 6 | 0 | 0 | 15 |
| 浸潤影 | 0 | 0 | 3 | 7 | 3 | 0 | 0 | 13 |
| 縦隔・肺門増大 | 0 | 0 | 1 | 1 | 1 | 0 | 0 | 3 |
| 腫瘤影 | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 1 |
| 胸水 | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 1 |
| 気胸 | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 1 |
| 受診者数(人) | 0 | 1,174 | 2,387 | 1,912 | 884 | 50 | 0 | 6,407 |