

## 8. 消化器

### (2) 上部消化管内視鏡検査

<検査結果判定> ※有所見率…B2 以上

①1日人間ドック

<男性>

	~29	30~39	40~49	50~59	60~69	70~79	80~	合計 (人)	比率 (%)
A	7	101	190	76	33	4	2	413	5.4
B1	12	324	848	682	388	69	8	2,331	30.6
B2	2	257	846	1,319	1,414	426	75	4,339	57.0
C	0	0	3	11	11	1	1	27	0.4
D	3	16	47	87	104	31	4	292	3.8
E	1	23	62	59	47	13	2	207	2.7
受診者数(人)	25	721	1,996	2,234	1,997	544	92	7,609	100.0
有所見者率(%)	24.0	41.1	48.0	66.1	78.9	86.6	89.1	63.9	

<女性>

	~29	30~39	40~49	50~59	60~69	70~79	80~	合計 (人)	比率 (%)
A	16	177	273	145	38	12	0	661	10.1
B1	11	342	988	849	429	83	10	2,712	41.5
B2	9	157	583	937	928	291	29	2,934	44.9
C	0	0	4	6	4	1	1	16	0.2
D	0	14	23	45	47	13	3	145	2.2
E	0	4	18	24	19	6	2	73	1.1
受診者数(人)	36	694	1,889	2,006	1,465	406	45	6,541	100.0
有所見者率(%)	25.0	25.2	33.2	50.4	68.1	76.6	77.8	48.4	

## ②2日人間ドック

&lt;男性&gt;

	~29	30~39	40~49	50~59	60~69	70~79	80~	合計 (人)	比率 (%)
A	3	47	58	44	19	6	2	179	4.9
B1	5	120	280	358	223	65	15	1,066	29.2
B2	1	73	270	617	667	399	117	2,144	58.7
C	0	0	4	5	4	1	0	14	0.4
D	0	5	17	48	45	20	20	155	4.2
E	0	4	15	30	28	20	0	97	2.7
受診者数(人)	9	249	644	1,102	986	511	154	3,655	100.0
有所見者率(%)	11.1	32.9	47.5	63.5	75.5	86.1	89.0	65.9	

&lt;女性&gt;

	~29	30~39	40~49	50~59	60~69	70~79	80~	合計 (人)	比率 (%)
A	1	28	33	26	24	7	1	120	8.1
B1	1	32	89	156	134	74	13	499	33.8
B2	1	16	55	178	315	177	37	779	52.7
C	0	1	0	1	6	5	1	14	0.9
D	0	0	5	10	12	7	6	40	2.7
E	0	0	3	2	10	7	3	25	1.7
受診者数(人)	3	77	185	373	501	277	61	1,477	100.0
有所見者率(%)	33.3	22.1	34.1	51.2	68.5	70.8	77.0	58.1	

## ③全国健康保険協会管掌健康保険生活習慣病予防健診 一般健康診査

&lt;男性&gt;

	~29	30~39	40~49	50~59	60~69	70~79	合計(人)	比率(%)
A	0	3	4	3	0	0	10	4.5
B1	0	14	39	23	7	0	83	37.6
B2	0	9	32	44	30	2	117	52.9
D	0	1	1	3	2	0	7	3.2
E	0	1	1	1	1	0	4	1.8
受診者数(人)	0	28	77	74	40	2	221	100.0
有所見者率(%)	0.0	39.3	44.2	64.9	82.5	100.0	57.9	

&lt;女性&gt;

	~29	30~39	40~49	50~59	60~69	70~79	合計(人)	比率(%)
A	2	4	15	6	2	0	29	11.3
B1	2	17	54	39	8	0	120	46.7
B2	0	17	26	47	8	3	101	39.3
C	0	0	0	0	1	0	1	0.4
D	0	0	2	4	0	0	6	2.3
受診者数(人)	4	38	97	96	19	3	257	100.0
有所見者率(%)	0.0	44.7	28.9	53.1	47.4	100.0	42.0	

## ④全国健康保険協会管掌健康保険生活習慣病予防健診 2次検査

&lt;男性&gt;

	40~49	50~59	60~69	合計(人)	比率(%)
B2	0	1	1	2	40.0
D	1	1	0	2	40.0
E	1	0	0	1	20.0
受診者数(人)	2	2	1	5	100.0
有所見者率(%)	100.0	100.0	100.0	100.0	

&lt;女性&gt;

	40~49	50~59	合計(人)	比率(%)
A	1	0	1	25.0
B1	1	0	1	25.0
D	0	2	2	50.0
受診者数(人)	2	2	4	100.0
有所見者率(%)	0.0	100.0	50.0	

## ⑤保険診療外来

&lt;男性&gt;

	30～39	40～49	50～59	60～69	70～79	80～	合計 (人)	比率 (%)
A	0	0	0	0	0	0	0	0.0
B1	4	2	7	4	3	0	20	11.2
B2	4	9	25	31	18	6	93	52.2
C	0	2	1	4	1	0	8	4.5
D	4	10	12	17	7	1	51	28.7
E	0	2	4	0	0	0	6	3.4
受診者数(人)	12	25	49	56	29	7	178	100.0
有所見者率(%)	66.7	92.0	85.7	92.9	89.7	100.0	88.8	

&lt;女性&gt;

	30～39	40～49	50～59	60～69	70～79	80～	合計 (人)	比率 (%)
A	0	0	0	0	0	0	0	0.0
B1	2	3	2	6	2	0	15	16.1
B2	1	8	21	16	7	0	53	57.0
C	0	0	1	0	0	0	1	1.1
D	0	4	6	11	2	1	24	25.8
E	0	0	0	0	0	0	0	0.0
受診者数(人)	3	15	30	33	11	1	93	100.0
有所見者率(%)	33.3	80.0	93.3	81.8	81.8	100.0	83.9	

## &lt;検査所見&gt;

①1日ドック

## &lt;男性&gt;

	~29	30~39	40~49	50~59	60~69	70~79	80~	合計(人)
異常なし	7	101	190	76	33	4	2	413
食道ポリープ	0	8	17	22	18	5	2	72
食道隆起性病変	1	1	4	1	2	1	0	10
食道陥凹性病変	0	1	2	2	4	1	0	10
食道粘膜下腫瘍	0	5	19	28	24	8	1	85
食道炎	8	210	593	702	593	124	24	2,254
食道潰瘍	0	0	0	1	0	1	0	2
急性胃炎	8	289	905	1,090	927	212	36	3,467
慢性胃炎	2	175	573	1,027	1,246	401	68	3,492
胃潰瘍	0	2	5	11	11	4	0	33
胃潰瘍瘢痕	0	2	33	76	124	36	7	278
胃ポリープ	6	124	464	638	473	133	27	1,865
胃隆起性病変	1	2	6	6	23	6	2	46
胃陥凹性病変	1	5	12	22	35	13	1	89
胃粘膜下腫瘍	0	10	45	84	109	27	8	283
十二指腸炎	0	35	105	117	68	16	1	342
十二指腸潰瘍	2	5	8	13	7	0	1	36
十二指腸潰瘍瘢痕	0	37	107	150	141	37	7	479
十二指腸ポリープ	0	1	21	41	65	11	3	142
十二指腸隆起性病変	0	1	9	10	10	2	0	32
十二指腸陥凹性病変	0	1	1	1	3	1	0	7
十二指腸粘膜下腫瘍	0	1	10	21	14	6	1	53
ブルンネル腺過形成	0	1	32	64	70	14	5	186
その他の病変	12	389	1,243	1,530	1,340	431	73	5,018
総計	48	1,406	4,404	5,733	5,340	1,494	269	18,694

&lt;女性&gt;

	~29	30~39	40~49	50~59	60~69	70~79	80~	合計(人)
異常なし	16	177	273	145	38	12	0	661
食道ポリープ	0	10	44	31	30	5	1	121
食道隆起性病変	0	0	2	3	4	0	0	9
食道陥凹性病変	0	0	0	1	1	0	0	2
食道粘膜下腫瘍	0	1	4	6	10	6	0	27
食道炎	1	64	204	249	216	76	5	815
急性胃炎	11	226	730	818	565	149	17	2,516
慢性胃炎	5	109	423	765	820	253	32	2,407
胃潰瘍	0	1	0	3	3	2	3	12
胃潰瘍瘢痕	0	1	15	28	38	13	0	95
胃ポリープ	4	226	779	814	502	132	16	2,473
胃隆起性病変	0	1	5	10	12	2	2	32
胃陥凹性病変	0	7	5	18	18	4	0	52
胃粘膜下腫瘍	1	13	64	80	84	34	4	280
十二指腸炎	0	4	19	22	14	8	1	68
十二指腸潰瘍	0	2	9	6	1	1	0	19
十二指腸潰瘍瘢痕	0	9	24	84	81	16	1	215
十二指腸ポリープ	0	0	8	13	14	1	0	36
十二指腸隆起性病変	0	0	2	3	2	0	0	7
十二指腸陥凹性病変	0	0	0	0	1	0	0	1
十二指腸粘膜下腫瘍	0	1	6	9	7	2	0	25
ブルンネル腺過形成	0	1	11	15	11	5	1	44
その他の病変	5	178	631	737	646	240	30	2,467
総計	43	1,031	3,258	3,860	3,118	961	113	12,384

## ②2日ドック

&lt;男性&gt;

	~29	30~39	40~49	50~59	60~69	70~79	80~	合計(人)
異常なし	3	47	58	44	19	6	2	179
食道ポリープ	0	1	5	10	14	5	2	37
食道隆起性病変	0	0	0	1	2	1	0	4
食道陥凹性病変	0	0	1	2	0	3	1	7
食道粘膜下腫瘍	0	0	3	14	9	13	1	40
食道炎	2	69	213	356	259	128	17	1,044
急性胃炎	1	104	285	539	440	199	45	1,613
慢性胃炎	1	51	186	467	597	365	138	1,805
胃潰瘍	0	0	2	4	7	2	2	17
胃潰瘍瘢痕	1	0	6	26	45	34	15	127
胃ポリープ	1	31	146	293	237	116	31	855
胃隆起性病変	0	0	1	8	9	6	6	30
胃陥凹性病変	0	2	5	12	16	6	8	49
胃粘膜下腫瘍	0	3	14	42	60	35	14	168
十二指腸炎	0	6	30	41	21	3	2	103
十二指腸潰瘍	0	2	0	7	2	2	0	13
十二指腸潰瘍瘢痕	0	6	28	81	80	37	17	249
十二指腸ポリープ	0	1	8	26	20	6	3	64
十二指腸隆起性病変	0	0	3	3	7	1	0	14
十二指腸粘膜下腫瘍	0	1	3	4	9	5	1	23
ブルンネル腺過形成	0	2	10	30	31	8	5	86
その他の病変	5	129	426	752	671	391	120	2,494
総計	14	455	1,433	2,762	2,555	1,372	430	9,021

&lt;女性&gt;

	~29	30~39	40~49	50~59	60~69	70~79	80~	合計(人)
異常なし	1	28	33	26	24	7	1	120
食道ポリープ	0	0	2	8	9	4	1	24
食道隆起性病変	0	0	0	1	2	3	0	6
食道陥凹性病変	0	0	0	0	0	1	1	2
食道粘膜下腫瘍	0	0	3	2	4	4	0	13
食道炎	1	8	17	61	85	55	14	241
急性胃炎	1	22	64	153	186	130	25	581
慢性胃炎	0	14	34	142	265	157	32	644
胃潰瘍	0	0	0	1	0	1	1	3
胃潰瘍瘢痕	0	1	0	2	7	4	7	21
胃ポリープ	0	16	70	139	146	90	21	482
胃隆起性病変	0	0	1	2	2	0	1	6
胃陥凹性病変	0	0	2	4	6	3	2	17
胃粘膜下腫瘍	0	3	9	24	28	21	7	92
十二指腸炎	0	0	3	5	4	1	0	13
十二指腸潰瘍	0	0	2	0	1	0	0	3
十二指腸潰瘍瘢痕	0	2	1	16	20	12	2	53
十二指腸ポリープ	0	0	0	3	5	2	0	10
十二指腸隆起性病変	0	0	0	0	1	1	0	2
十二指腸粘膜下腫瘍	0	0	2	1	3	0	0	6
ブルンネル腺過形成	0	0	1	1	5	2	0	9
その他の病変	0	13	53	145	201	139	41	592
総計	3	107	297	736	1,004	637	156	2,940

## ③全国健康保険協会管掌健康保険生活習慣病予防健診 一般健康診査

&lt;男性&gt;

	~29	30~39	40~49	50~59	60~69	70~79	合計（人）
異常なし	0	3	4	3	0	0	10
食道ポリープ	0	3	0	1	1	0	5
食道粘膜下腫瘍	0	0	0	2	0	0	2
食道炎	0	3	21	20	8	0	52
急性胃炎	0	13	35	27	23	0	98
慢性胃炎	0	6	17	37	27	1	88
胃潰瘍	0	0	1	2	0	0	3
胃潰瘍瘢痕	0	0	1	4	1	0	6
胃ポリープ	0	5	25	19	10	0	59
胃隆起性病変	0	0	0	1	0	0	1
胃陥凹性病変	0	0	1	0	0	0	1
胃粘膜下腫瘍	0	0	4	4	3	0	11
十二指腸炎	0	1	5	3	2	0	11
十二指腸潰瘍	0	1	0	0	0	0	1
十二指腸潰瘍瘢痕	0	0	3	6	6	0	15
十二指腸隆起性病変	0	1	0	0	0	0	1
十二指腸粘膜下腫瘍	0	0	0	2	0	0	2
ブルンネル腺過形成	0	0	0	1	0	0	1
その他の病変	0	20	54	46	26	4	150
総計	0	56	171	178	107	5	517

## &lt;女性&gt;

	~29	30~39	40~49	50~59	60~69	70~79	合計(人)
異常なし	2	4	15	6	2	0	29
食道ポリープ	0	1	2	0	0	0	3
食道粘膜下腫瘍	0	0	0	1	0	0	1
食道炎	0	3	11	9	4	1	28
急性胃炎	2	13	36	33	8	1	93
慢性胃炎	0	10	20	39	8	3	80
胃潰瘍瘢痕	0	1	0	4	0	1	6
胃ポリープ	0	17	40	36	8	0	101
胃陥凹性病変	0	0	1	3	0	0	4
胃粘膜下腫瘍	0	1	1	5	0	0	7
十二指腸炎	0	0	0	2	0	0	2
十二指腸潰瘍瘢痕	0	2	1	7	0	0	10
十二指腸粘膜下腫瘍	0	0	1	0	0	0	1
ブルンネル腺過形成	0	0	1	0	0	0	1
その他の病変	1	10	31	35	10	2	89
総計	5	62	160	180	40	8	455

④全国健康保険協会管掌健康保険生活習慣病予防健診 2次検査

## &lt;男性&gt;

	40~49	50~59	60~69	合計(人)
食道炎	2	0	0	2
急性胃炎	2	0	0	2
慢性胃炎	0	4	1	5
胃潰瘍	0	1	0	1
十二指腸炎	1	0	0	1
十二指腸潰瘍	1	0	0	1
その他の病変	3	0	1	4
総計	9	5	2	16

## &lt;女性&gt;

	40~49	50~59	合計(人)
異常なし	1	0	1
急性胃炎	2	2	4
慢性胃炎	0	2	2
胃潰瘍瘢痕	0	1	1
胃ポリープ	1	0	1
胃陥凹性病変	0	1	1
十二指腸潰瘍瘢痕	0	1	1
その他の病変	1	0	1
総計	5	7	12

## ⑤保険診療外来

&lt;男性&gt;

	30～39	40～49	50～59	60～69	70～79	80～	合計(人)
食道ポリープ	0	0	1	1	0	0	2
食道陥凹性病変	0	0	0	0	2	0	2
食道粘膜下腫瘍	0	0	0	1	0	0	1
食道炎	5	9	17	14	4	1	50
急性胃炎	9	13	30	27	11	1	91
慢性胃炎	5	21	37	51	30	6	150
胃潰瘍	0	3	2	1	0	0	6
胃潰瘍瘢痕	0	6	5	11	3	0	25
胃ポリープ	2	1	10	9	5	1	28
胃隆起性病変	0	0	0	5	0	0	5
胃陥凹性病変	1	2	6	8	4	1	22
胃粘膜下腫瘍	1	2	1	3	1	1	9
十二指腸炎	1	1	3	2	0	0	7
十二指腸潰瘍	0	3	0	0	0	0	3
十二指腸潰瘍瘢痕	0	3	5	5	2	0	15
十二指腸ポリープ	0	0	1	1	0	0	2
十二指腸隆起性病変	0	0	2	2	0	0	4
ブルンネル腺過形成	0	0	1	0	0	0	1
その他の病変	8	8	28	45	18	6	113
総計	32	72	149	186	80	17	536

&lt;女性&gt;

	30~39	40~49	50~59	60~69	70~79	80~	合計(人)
食道ポリープ	0	0	1	0	2	0	3
食道隆起性病変	0	0	0	1	0	0	1
食道粘膜下腫瘍	0	0	1	1	1	0	3
食道炎	0	2	4	4	4	0	14
急性胃炎	1	4	14	18	6	0	43
慢性胃炎	1	13	24	21	7	1	67
胃潰瘍	0	1	2	1	0	0	4
胃潰瘍瘢痕	0	0	2	1	0	0	3
胃ポリープ	1	3	7	13	5	1	30
胃隆起性病変	0	1	1	0	0	1	3
胃陥凹性病変	0	2	2	2	2	0	8
胃粘膜下腫瘍	0	0	1	4	1	0	6
十二指腸潰瘍瘢痕	0	1	0	0	0	0	1
ブルンネル腺過形成	0	1	0	0	0	0	1
その他の病変	0	4	11	22	5	1	43
総計	3	32	70	88	33	4	230

## &lt;胃組織検査結果&gt;

## &lt;男性&gt;

	~29	30~39	40~49	50~59	60~69	70~79	80~	合計 (人)	比率 (%)
Group I	3	18	57	112	111	33	13	347	86.3
Group II	0	0	1	4	7	2	1	15	3.7
Group III	0	0	1	0	5	3	0	9	2.2
Group IV	0	2	1	0	2	3	2	10	2.5
Group V	0	0	2	4	11	2	2	21	5.2
受診者数(人)	3	20	62	120	136	43	18	402	100.0

## &lt;女性&gt;

	30~39	40~49	50~59	60~69	70~79	80~	合計 (人)	比率 (%)
Group I	9	25	55	52	14	6	161	91.0
Group II	0	0	0	2	0	0	2	1.1
Group III	0	0	1	1	2	0	4	2.3
Group IV	0	0	1	0	1	0	2	1.1
Group V	0	1	3	3	0	0	7	4.0
その他	0	0	0	1	0	0	1	0.6
受診者数(人)	9	26	60	59	17	6	177	100.0

Group I : 正常組織および異型を示さない良性病変

Group II : 異型を示すが良性と判定される病変

Group III : 良性と悪性の境界領域の病変

Group IV : 癌が強く疑われる病変

Group V : 癌

その他 : 検体不適